

Westlake Plastic Surgery™

DR. ROBERT CARIDI

Aspirin Use

The following is a partial list of medications that can increase your bleeding tendency at the time of surgery and in the immediate post operative period.

This is a **PARTIAL** list. Any and all medications (including herbal products and vitamins) that you are taking must be reported to our office a minimum of **6 weeks prior to your surgery**. If you are unsure about a product you are using that is not on this list, call our office and ask the nursing staff. Check the labels of ALL medications that you take for ingredients that may be on this list.

DO NOT take any drugs on this list **6 weeks (3 weeks for any Diet Aids)** prior to or following your surgery without Dr. Caridi's consent. **If you are unsure of any product you wish to use, please call the office before taking it or use Tylenol.**

Aspirin	Buf Tabs	Emprin	Oxycodone	Voltaren
Actifed	Brufen	Emprazil	Pabirin	Wesprin
Advil	Bupropion Cama	Excedrin	Panalgesic	Zactin
Aleve	Cephalgesic	Feldene	Pepto Bismol	Zorpin
Allegra D	Claritin D	Fiorinal	Percodan	Zyrtec D
Alka Seltzer	Clinoril	Flurbiprophen	Persisrin	
Aluprin	Congespirin	Sodium	Quagesic	
Alcohol	Cope	Four Way	Rhinex	
Anacin	Coricidan D	Cold Tabs	Robaxisal	
Anaprox	Coricidin Tablets	Ibuprofen	Robaxin	
Anaproxn	Contact	Indocin	Rufen	
Ansaid	Coumadin	Indomethacin	Sine Off	
APA	Concerta Darvon	Lodinc	Sine Aid	
Arthritis Pain	with Aspirin	Meclomen	Sudafed	
Reliever	Diet Aids	Medipren	Supac	
ASA	Decongestants	Midol	Synalogos	
Asperbuf	Diurex	Mobigesic	Talwin	
Asriptin	Doan's Pills	Monacet	Toradol	
Aspergum	Dolobid	Motrin	Triaminiein	
BC Powder	Dirstan	Norgesic	Trandate	
Bayer	Duradyne	Nardil	Trental	
Bufferin	Easprin	Nuprin	Trilisate	
Buffex	Ecotrin	Nyquil	Vanquish	